
Proportion of HIV/AIDS Cases among Adults and Adolescents, by Sex and Age Group Diagnosed in 2004—33 States





Note. Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis Data from 33 states with confidential name-based HIV infection reporting since at least 2000. Data based on person's age at diagnosis. Data have been adjusted for reporting delays.



HIV/AIDS Cases among Male Adolescents and Young Adults, by Transmission Category 2001–2004—33 States

Transmission category	13-19	years	20-24 y N	years
		,		
Male-to-male sexual contact	1,723	76	6,485	74
Injection drug use (IDU)	185	8	734	8
Male-to-male sexual contact				
and IDU	85	4	448	5
High risk heterosexual contact*	278	12	1,121	13
Other/not identified**	11	<1	32	<1
Total	2,282	100	8,820	100

Note. Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2000. Data have been adjusted for reporting delays.



^{*} Heterosexual contact with a person known to have or at high risk for HIV infection.



^{**} Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

HIV/AIDS Cases among Female Adolescents and Young Adults, by Transmission Category 2001–2004—33 States

T	13-19 years		20-24	20-24 years	
Transmission Category	N	%	N		
Injection drug use	285	14	697	15	
High risk heterosexual contact*	1,739	85	3,929	84	
Other/not Identified**	22	1	46	1	
Total	2,046	100	4,672	100	

Note. Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2000. Data have been adjusted for reporting delays.






^{*} Heterosexual contact with a person known to have or at high risk for HIV infection.

^{**} Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

HIV/AIDS Cases among Adolescents 13 to 19 Years of Age, 2004—33 States






Note. Data include persons with a diagnosis of HIV infection regardless of AIDS status of diagnosis. Data from 33 states with confidential name-based infection reporting since at least 2000. Data have been adjusted for reporting delays. Data based on person's age at diagnosis.




Adolescents 13 to 19 Years of Age Living with HIV/AIDS, 2004—33 States








Proportion of HIV/AIDS Cases and Population among Adolescents 13 to 19 Years of Age, by Race/Ethnicity Diagnosed in 2004—33 States







Proportion of HIV/AIDS Cases and Population among Young Adults 20 to 24 Years of Age, by Race/Ethnicity Diagnosed in 2004—33 States





Note. Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2000. Data have been adjusted for reporting delays.

* Include 23 persons of unknown race or multiple races.



Proportion of AIDS Cases among Adults and Adolescents, by Sex and Age Group, Reported in 2004—United States







AIDS Cases among Male Adolescents and Young Adults, by Transmission Category 2000–2004—United States

Transmission category	13-19 N	years %	20-24 ₎ N	ears %
Male-to-male sexual contact	564	58	3,314	69
Injection drug use (IDU)	108	11	511	11
Male-to-male sexual contact and IDU	35	4	263	5
High risk heterosexual contact*	123	13	688	14
Other/not identified**	149	15	62	1
Total	979	100	4,838	100

Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.





^{*} Heterosexual contact with a person known to have or at high risk for HIV infection.
** Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

AIDS Cases among Female Adolescents and Young Adults, by Transmission Category 2000–2004—United States

	13-19 years	20-24 years	
Transmission Category	<u>N</u> %	<u>N</u>	<u></u> %
Injection drug use	115 13	480	17
High risk heterosexual contact*	548 64	2,223	81
Other/not Identified**	189 22	51	2
Total	852 100	2,754	100

Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.






^{*} Heterosexual contact with a person known to have or at high risk for HIV infection.

^{**} Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.


Reported AIDS Cases among Adolescents 13 to 19 Years of Age, by Sex,1985–2004—United States N=5,593





Note. Data based on person's age at diagnosis.

Proportion of AIDS Cases and Population among Adolescents 13 to 19 Years of Age, by Race/Ethnicity Reported in 2004—50 States and DC






Note. Data based on person's age at AIDS diagnosis. Excludes persons from US dependencies, possessions, and associated nations.

*Includes 1 person of unknown race or multiple races.




Reported AIDS Cases among Adolescents 13 to 19 Years of Age, 2004—United States






Adolescents 13 to 19 Years of Age Reported to be Living with AIDS, 2004—United States






Reported AIDS Cases among Young Adults 20 to 24 Years of Age, by Sex,1985–2004—United States N=32,757





Note. Data based on person's age at diagnosis.

Proportion of AIDS Cases and Population among Young Adults 20 to 24 Years of Age, by Race/Ethnicity Reported in 2004—50 States and DC





Note. Data based on person's age at AIDS diagnosis. Excludes persons from US dependencies, possessions, and associated nations. * Includes 10 persons of unknown race or multiple races.

